

Post Operative Info - Breast Conservation Surgery

Post-Breast Conservation Surgery advice

Care of your wound

- Your wound will be covered with a waterproof dressing. This will still be intact until your post-surgery appointment.
- You may notice some dry blood under the dressing. This is very normal and is actually helping your wound heal.
- You do not need to do anything special to your wound whilst the dressings are in place. After the dressings are removed Dr. Green will advise you of various treatments to decrease scarring.

Stitches

- Your wound has been closed entirely with dissolving sutures. These will take several weeks to completely dissolve. You will not need to have any stitches removed after your surgery.

Drains

- Drains are rarely placed for breast conservation surgery.
- If you have also undergone an Axillary dissection at the same time then a drain will be placed for this operation. Continue reading below in the '*axillary node surgery section*' for advice about axillary drains.

Signs of Infections

- Bad Infection is a rare complication after a breast surgery. However, it is possible for a small infection to occur that can be easily treated with tablet antibiotics.
- It is normal for the wound itself to look red. However if the redness is increasing past your dressings you may have an early infection.

- If you are concerned notify Dr. Green or contact your GP who may commence you on a course of tablet antibiotics.

Pain Relief

- Breast surgery is not usually very painful. Your breast will feel heavy and swollen for a week or two. Most people can manage with simple analgesia (paracetamol) and anti-inflammatory (Nurofen). Some people do require some stronger pain relief and this will be required if necessary prior to your discharge.
- A supportive Bra (such as a sports bra can provide some relief from the heavy feeling
- Avoid constipation whilst taking the pain relief. You should see your Pharmacist for some coloxyl and senna or Movichol whilst taking pain relief.

Showering and deodorant

- Your dressings are waterproof and it is perfectly safe for you to have a shower after your surgery. When you have finished your shower just pat your wound dry.
- Deodorant (aerosol deodorant or roll on) is perfectly safe to use after you surgery. It is best to avoid powdered based deodorants.

Return to work

- You will need to have some time off work after your surgery.
- Dr. Green will happily provide a medical certificate for these times if you wish.

Energy levels and resuming normal activities

- It is normal to not have full energy levels after the operation.
- You should be able to perform usual light duties within 1 week of the surgery. Although it is quite normal to not regain full levels of energy for several weeks after an anaesthetic.
- Regular light exercise (walking, riding etc..) is encouraged after your surgery. A basic rule is 'if it hurts its to strenuous and you should stop'
- Regular exercise is important. There is good evidence supporting exercise and healthy eating in terms of your breast cancer treatment.

Follow Up

Dr. Green will normally see you in the Rooms 1 week after your operation. Please call 32263800 to confirm an appointment.

Information specifically for Axillary Lymph Node Surgery

Sentinel Node Biopsy

- The blue dye used to find the node will make your Urine Blue for a day. This is normal and not dangerous. It will also give you a grey complexion for 24 hours but will pass.
- No additional post surgery care is required.
- There is NO NEED to avoid blood tests or blood pressure taking from your arm after a sentinel node biopsy.
- The very small risk of lymphedema will be monitored by Dr. Green.

Axillary Clearance

- **Drains**
 - You will have a drain in place after and axillary clearance surgery. This is important to remove fluid that is after surgery
 - The nursing staff will teach you how to look after the drain and provide you with spares drainage bags and a carry bag. Any problems with call the Ward directly (any time) or Dr. Greens rooms (in business hours).
 - You will be shown how to record how much is coming into the drain. It will initially be a large amount of red/yellow fluid. Once it is less than 50ml on two consecutive days the drain will be remove. On average this takes about 1 week, although in some cases it may be 2-3 weeks.
 - If you accidentally pull your drain out DO NOT PANIC. Nothing bad will occur. Let Dr. Green know and he will arrange for the fluid that builds up to be drained.
- An axillary clearance is often more uncomfortable and you will be given stronger pain relief. It is important to avoid becoming constipated while on these medications.
- Physiotherapy will be started whilst you are in hospital. You will be given a number of exercises to do. It is STRONGLY encouraged to continue these exercises whilst at home. It will help in the prevention of lymphedema. If necessary additional physiotherapy can be arrange when you return for your post surgery appointment.
- Shoulder movements. It is perfectly safe to raise your arms above your head. You should not restrict any movements.
- Lymphedema will not develop overnight. Early swelling is most likely secondary to the surgery and fluid you where given. Dr. Green will monitor you for signs of lymphedema and if it begins will arrange expert lymphedema care immediately.

It is best to avoid blood tests or blood pressure taking from your arm after an axillary clearance until you have been cleared of any lymphoedema.